



“Gheorghe Asachi” Technical University of Iasi, Romania



STUDY OF THE DEVOLATILIZATION KINETICS OF KENAF BY ISOCONVERSIONAL METHODS. INFLUENCE OF VARIABLES

S. Román^{1*}, B. Ledesma¹, A. Álvarez-Murillo¹, A. Al-Kassir², J.F. González¹

¹Department of Applied Physics

²Mechanics, Energetic and Materials Department, University of Extremadura, Avda de Elvas s/n, Badajoz 06006, Spain

Abstract

The thermal degradation of kenaf (*Hibiscus cannabinus*) under the effect of several variables (particle size, Ar flow rate, sample initial mass and heating rate) was investigated. From experimental data and by application of various model-free methods (Friedman, Kissinger-Akahira-Sunose and Flynn-Wall-Ozawa), the activation energies (E_A) were calculated as a function of conversion. The results obtained allowed to conclude that kenaf follows a multi-stage degradation mechanism, with E_A values ranging 170-225 kJ mol⁻¹. Despite the slight differences showed by the three methods on the E_A values, they agreed about the pattern of E_A with conversion.

Key words: kenaf, kinetics, pyrolysis, model-free methods

Received: May, 2013; *Revised final:* August, 2014; *Accepted:* August, 2014

*Author to whom all correspondence should be addressed: e-mail: sroman@unex.es, Phone: +34 924289600, Fax: +34 924289601