



"Gheorghe Asachi" Technical University of Iasi, Romania



MONITORING THE BREAKDOWN OF DINOCAPI IN SPIKED SOIL WINE AND GRAPE SAMPLES BY GC/MS AND FTIR/ATR

Nicoleta Grigoriu¹, Catalina Calin^{2*}, Gina Vasile Scaeteanu³, Roxana Maria Madjar³
Rodica Mihaela Lungu¹, Gabriel Epure¹

¹Scientific Research Center for CBRN Defense and Ecology, Oltenitei Av. 225, Bucharest, Romania

²Petroleum-Gas University of Ploiesti, Bucharest Av. 39, Ploiesti, Romania

³University of Agronomical Sciences and Veterinary Medicine, Marasti 59 Str., Bucharest, Romania

Abstract

Capillary gas chromatography coupled with mass spectrometry technique was used for the assessment of low concentration of dinitrophenol pesticide, Dinocap in different spiked samples. For this purpose it has been spiked with Dinocap (15- 200 µg/mL) five environmental matrices (a soil sample and two each samples of wine and grapes). The Dinocap degradation grade in these samples was also monitored. The extraction was performed in dichloromethane. The assessment of the Dinocap in the samples were also achieved by FTIR/ATR and reveals that after 18 days of pesticide application the recovery level of Dinocap decreased from 64% to 3% in grape, from 72% to 9% in soil, from 82% to 10% in wine at 18 days after pesticide application. The method described in this paper could be used for screening and identification of the named pesticide in other environmental samples with a spiking level as low as 2-20 µg/mL.

Key words: dinitrophenol pesticide, dinocap, FTIR, GC/MS, wine

Received: November, 2013; *Revised final:* January, 2015; *Accepted:* January, 2015

*Author to whom all correspondence should be addressed: e-mail: catalina.calin20@yahoo.com; Phone: 0722738775