Environmental Engineering and Management Journal

August 2021, Vol. 20, No. 8, 1309-1315 http://www.eemj.icpm.tuiasi.ro/; http://www.eemj.eu



"Gheorghe Asachi" Technical University of lasi, Romania



CHARACTERIZATION AND INVESTIGATION OF CHICKEN LITTER AS A SOURCE OF PHOSPHORUS FOR FERTILIZERS

Carla Oliveira Nascimento^{1*}, Sulian Junkes Dal Molin², Vinícius de Melo Benites³, Elaine Christine de Magalhães Cabral-Albuquerque¹, Rosana Lopes Fialho¹

¹Department of Industrial Engineering, Federal University of Bahia, 02 Prof. Aristides Novis Street, Federação, Salvador, BA, Brazil

²Department of Soil Science, Santa Catarina State University, 2090 Luiz de Camões Avenue, Conta Dinheiro, Lages, SC, Brazil ³Brazilian Agricultural Research Corporation – Soils, 1024 Jardim Botânico Street, Jardim Botânico, Rio de Janeiro, RJ, Brazil

Abstract

Chicken production has been growing in Brazil due to population growth and consequently the waste generated in this process, namely chicken litter, has also increased. Currently, the main destination of chicken litter is its application to the soil in areas near to the chicken production units. The objective of this work was to characterize the chicken litter produced in the Brazilian Midwest using different techniques, in order to diversify the application of this residue and minimize the environmental impacts generated by chicken meat production. Chicken litter characterization was performed by analyses of particle size distribution, moisture content, pH, water activity, proximate composition, total nitrogen, protein and non-protein nitrogen, and pyrolysis and thermogravimetric analysis. In addition, a greenhouse experiment was carried out to evaluate the agronomic efficiency of chicken litter compared to mineral fertilizer. Mass balance of chicken litter pyrolysis showed that it kept 50% solid mass and the first most significant mass loss of chicken litter occurs between 250 and 350 °C from the thermogravimetric analysis. Chicken litter was less efficient than mineral fertilizer. However, the great improvement in shoot dry matter yield compared to control showed the potentiality of this residue as a nutrient source when properly used.

Key words: nutrient resource, organic fertilizer, plant nutrition, poultry waste

Received: March, 2020; Revised final: January, 2021; Accepted: March, 2021; Published in final edited form: August, 2021

^{*} Author to whom all correspondence should be addressed: e-mail: carla.nascimento@ufba.br