



---

## WATER, VEHICLE FOR PREVENTIVE ORAL HEALTH PRODUCTS

**Mihaela Raescu<sup>1\*</sup>, Carmen Defta<sup>2</sup>, Gabriela Bancescu<sup>2</sup>, Mihaela Tuculina<sup>3</sup>,  
Marian Vladimir Constantinescu<sup>4</sup>**

<sup>1</sup>*"Titu Maiorescu" University, Faculty of Medicine and Dentistry, Department of Preventive Dentistry, 67A Gheorghe Petrașcu Street, 3 District, 031593 Bucharest, Romania*

<sup>2</sup>*"Carol Davila" University of Medicine and Pharmacy, Faculty of Dentistry, Department of Microbiology, 19 Plevan Avenue, Bucharest, Romania*

<sup>3</sup>*University of Medicine and Pharmacy, Faculty of Dentistry, Department of Restorative Odontototerapy, 2 Petru Rareș Street, 200349 Craiova, Romania*

<sup>4</sup>*"Titu Maiorescu" University, Faculty of Medicine and Dentistry, Department of Oral Rehabilitation, 67A Gheorghe Petrașcu Street, 3 District, 031593 Bucharest, Romania*

---

### Abstract

This study highlights the effectiveness of mouthwash compared with the gel Dentomer on some microbial strains of oral streptococci: mitis, mutans and oralis. Mitis and mutans strains of oral streptococci are sensitive to mouthwash DENTOMER and insensitive to the DENTOMER gel, while St oralis is resistant to both antiseptics that were tested. Dentomer mouthwash can be used as a preventive method, being effective against plaque accumulation.

*Key words:* mouthwash, oral streptococci, prevention

*Received: September, 2010; Revised: November, 2010; Accepted: November, 2010*

---

---

\* Author to whom all correspondence should be addressed: e-mail: [mihaela.raescu@gmail.com](mailto:mihaela.raescu@gmail.com); Phone: +40 744 704088